

## Recommendation

This study aimed to investigate the transcriptome-wide screening of expression of genes, lncRNAs and circRNAs in response to *H. theivora* infestation in the *C. sinensis* plant. Biochemical analysis involving estimation of total phenolics and flavonoids, antioxidative enzymatic assays of five different enzymes viz. POX, APX, PPO, PAL and CAT, revealed a significant change in biochemical activities in the plant in response to the TMB feeding. The RNA-seq analysis involving three replicates of control and three of infested, provided a detailed description of the expression pattern of the transcripts in the tea plant tissues. Accordingly, functional annotation analysis of the transcripts indicated how they might have played significant roles in response of the plant to the feeding insect.

In the present study, 9502 lncRNAs were identified from healthy and TMB-infested *C. sinensis* tissues using high-throughput RNA sequencing data, out of which 80 lncRNAs got differentially expressed in response to TMB infestation. Determination of genes that could act as potential targets of lncRNAs revealed that the identified lncRNAs could possibly target as many as 5804 genes. Differential gene expression (DGE) analysis led to the identification of 3665 differentially expressed genes (DEGs), of which, expression of 1767 genes got upregulated and 1898 genes got downregulated during tea plant's response to TMB. Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis of DEGs and lncRNA-target genes have shown that TMB infestation can trigger transcriptomic reprogramming possibly leading to altered primary and secondary metabolism in *C. sinensis*. Identification of TMB-responsive circRNAs (another class of non-coding RNA) in tea plant revealed differential expression of 34 circRNAs during TMB feeding. These circRNAs are found

to be associated with biological pathways like “tryptophan metabolism”, “biosynthesis of secondary metabolites”, “porphyrin metabolism”, “nucleocytoplasmic transport”, “N-glycan biosynthesis” etc. LncRNAs and circRNAs can act as competing endogenous RNAs (ceRNAs) to bind with common microRNA (miRNA) response elements (MREs) involving a competition between mRNAs, lncRNAs and circRNAs. Assessment of this “sponging” activity of lncRNAs and circRNAs led to the construction of the ceRNA network consisting of 5 lncRNAs, 17 circRNAs, 33 mRNAs competing against each other to bind with 37 miRNAs. The expression of 6 DEGs and 5 differentially expressed lncRNAs (DELs) have been validated by qRT-PCR. This study highlights the reprogramming in expression of coding and non-coding transcripts as a part of defense regulatory network in tea plants against TMB.

*Niraj Agarwal*  
(Dr. Niraj Agarwala)

Supervisor

*Kuntala Sarma Bordoloi*  
(Kuntala Sarma Bordoloi)

Research scholar